

Viticulture Foliar Crop Program

A foliar crop program suitable for wine and table grapes.

Analysis

	Copper as Cuprous Oxide	Zinc as Zinc Oxide	Manganese as Manganese Carbonate
Verno Copper + Zinc	300g/kg	300g/kg	-
Verno Manganese	-	-	340g/kg
Nordox 750WG Copper Fungicide	750g/kg	-	-

- Water dispersible granule formulation for free flowing ease of handling and low dust.
- Verno foliar fertilisers can be mixed in any ratio for the desired balance of copper, zinc and manganese.
- Verno Micronutrients and Nordox 750WG are certified for organic production.
- Verno foliar fertilisers are compatible with Nordox 750WG Copper fungicide and a wide range of insecticides, fungicides and foliar fertilisers when the spray solution is above pH 6.0.



Foliar applied fertiliser for the correction of micronutrient deficiencies.

Verno contains micronised particles which adhere strongly to plant surfaces, providing a reservoir of slow release micronutrients that are readily available to the plant.



Foliar Application Grape Vines

- Multiple low rate applications of Verno during pre flowering vegetative growth, post flowering and post harvest give the best results. The recommended concentration applies to a single Verno product or a combination of Verno products, depending on crop requirements.
- **Pre and Post Flowering:** Apply Verno at 100g - 500g/ha (maximum concentration 1g/1L). Multiple low rate applications are most effective.
- **Post Harvest:** Apply Verno at 500g/ha (maximum concentration 1g/1L).
- Nordox 750WG - For Downy Mildew control apply 100-130g/100L. First spray when shoots are 10cm long and re-apply at 10-14 day intervals while conditions are favourable for infection. (Refer to Nordox 750WG Fungicide label for more detailed instructions.)

Mixing Instructions

- Fill the spray tank to 3/4 with water, then add the required amount of Verno granules to the tank with continuous vigorous agitation.
 - Completely disperse the granules in the spray tank before adding any other mixing partners.
 - Continuous vigorous agitation is required to maintain suspension of the micronised particles.
 - Do not mix with other products prior to complete suspension. Mixing with other products is not recommended unless the compatibility is confirmed by the manufacturer of that product.
 - Completely flush tank and lines immediately after use.
-
- When applying mixtures of Verno products, the suggested rate and concentration applies to the total of the combined products. Care should be taken to not exceed the maximum rate and concentration for application at that growth stage.
 - Maximum 5g/L spray tank concentration.
 - Constant agitation is required to maintain Verno in suspension.



For more information

P +61 7 4639 2009
E office@tanuki.com.au

Tanuki Pty Ltd
27 Heinemann Road
Wellcamp QLD 4350
Australia

verno.com.au

While Tanuki Pty Ltd has taken all responsible care in the preparation of this technical sheet, the information is a guide only and should not be relied upon as a substitute for you seeking professional advice in relation to the use of Verno products. Verno and Nordox are a registered trademarks of Nordox AS.